



SP110 gene

SP110 nuclear body protein

Normal Function

The *SP110* gene provides instructions for making a protein called SP110 nuclear body protein, which is a component of cellular structures called nuclear bodies. Nuclear bodies are located within the nuclei of cells, where they help control the activity of certain genes. Nuclear bodies are also involved in the regulation of cell division, the self-destruction of cells that are damaged or no longer needed (apoptosis), and the normal function of the immune system.

SP110 nuclear body protein is active primarily in immune system cells called leukocytes and in the spleen. It likely helps regulate the activity of genes needed for the body's immune response to foreign invaders (such as viruses and bacteria).

Health Conditions Related to Genetic Changes

hepatic veno-occlusive disease with immunodeficiency

At least five mutations in the *SP110* gene have been found to cause hepatic veno-occlusive disease with immunodeficiency (VODI). Each of these mutations leads to the production of a nonfunctional version of SP110 nuclear body protein. A lack of functional protein impairs the immune system's ability to fight off foreign invaders, allowing recurrent and persistent infections to develop. It is unclear how the loss of SP110 nuclear body protein disrupts blood flow in the liver, leading to enlargement of the liver (hepatomegaly), a buildup of scar tissue (hepatic fibrosis), and liver failure.


other disorders

Several common variations (polymorphisms) in the *SP110* gene have been studied as risk factors for lung (pulmonary) tuberculosis. This disease is a contagious bacterial infection. At least one study has found that certain variations in the *SP110* gene may influence the risk of infection with the bacteria that cause pulmonary tuberculosis. However, several other studies have not found such an association. *SP110* variations are probably not a major genetic risk factor for this disease.

Chromosomal Location

Cytogenetic Location: 2q37.1, which is the long (q) arm of chromosome 2 at position 37.1

Molecular Location: base pairs 230,165,466 to 230,225,729 on chromosome 2 (Homo sapiens Annotation Release 108, GRCh38.p7) (NCBI)



Credit: Genome Decoration Page/NCBI

Other Names for This Gene

- FLJ22835
- IFI41
- IFI75
- interferon-induced protein 41, 30kD
- interferon-induced protein 41/75
- interferon-induced protein 75, 52kD
- IPR1
- nuclear body protein SP110
- phosphoprotein 41
- phosphoprotein 75
- SP110_HUMAN
- speckled, 110-KD
- transcriptional coactivator Sp110

Additional Information & Resources

Educational Resources

- Nuclear Protein Database: PML Nuclear Bodies
<http://npd.hgu.mrc.ac.uk/user/compartment?page=pml>

GeneReviews

- Hepatic Veno-Occlusive Disease with Immunodeficiency
<https://www.ncbi.nlm.nih.gov/books/NBK1271>

Scientific Articles on PubMed

- PubMed
<https://www.ncbi.nlm.nih.gov/pubmed?term=%28SP110%5BTIAB%5D%29+AND+%28%28Genes%5BMH%5D%29+OR+%28Genetic+Phenomena%5BMH%5D%29%29+AND+english%5Bla%5D+AND+human%5Bmh%5D+AND+%22last+3600+days%22%5Bdp%5D>

OMIM

- NUCLEAR BODY PROTEIN SP110
<http://omim.org/entry/604457>

Research Resources

- ClinVar
<https://www.ncbi.nlm.nih.gov/clinvar?term=SP110%5Bgene%5D>
- HGNC Gene Family: Minor histocompatibility antigens
<http://www.genenames.org/cgi-bin/genefamilies/set/870>
- HGNC Gene Family: PHD finger proteins
<http://www.genenames.org/cgi-bin/genefamilies/set/88>
- HGNC Gene Symbol Report
http://www.genenames.org/cgi-bin/gene_symbol_report?q=data/hgnc_data.php&hgnc_id=5401
- NCBI Gene
<https://www.ncbi.nlm.nih.gov/gene/3431>
- UniProt
<http://www.uniprot.org/uniprot/Q9HB58>

Sources for This Summary

- Babb C, Keet EH, van Helden PD, Hoal EG. SP110 polymorphisms are not associated with pulmonary tuberculosis in a South African population. Hum Genet. 2007 May;121(3-4):521-2. Epub 2007 Feb 8.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17287948>
- Bloch DB, Nakajima A, Gulick T, Chiche JD, Orth D, de La Monte SM, Bloch KD. Sp110 localizes to the PML-Sp100 nuclear body and may function as a nuclear hormone receptor transcriptional coactivator. Mol Cell Biol. 2000 Aug;20(16):6138-46.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/10913195>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC86089/>

- Roscioli T, Cliffe ST, Bloch DB, Bell CG, Mullan G, Taylor PJ, Sarris M, Wang J, Donald JA, Kirk EP, Ziegler JB, Salzer U, McDonald GB, Wong M, Lindeman R, Buckley MF. Mutations in the gene encoding the PML nuclear body protein Sp110 are associated with immunodeficiency and hepatic veno-occlusive disease. *Nat Genet.* 2006 Jun;38(6):620-2. Epub 2006 Apr 30.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16648851>
- Szeszko JS, Healy B, Stevens H, Balabanova Y, Drobniewski F, Todd JA, Nejntsev S. Resequencing and association analysis of the SP110 gene in adult pulmonary tuberculosis. *Hum Genet.* 2007 Apr;121(2):155-60. Epub 2006 Dec 6.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/17149599>
- Thye T, Browne EN, Chinbuah MA, Gyapong J, Osei I, Owusu-Dabo E, Niemann S, Rüscher-Gerdes S, Horstmann RD, Meyer CG. No associations of human pulmonary tuberculosis with Sp110 variants. *J Med Genet.* 2006 Jul;43(7):e32.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16816019>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC2564561/>
- Tosh K, Campbell SJ, Fielding K, Sillah J, Bah B, Gustafson P, Manneh K, Lisse I, Sirugo G, Bennett S, Aaby P, McAdam KP, Bah-Sow O, Lienhardt C, Kramnik I, Hill AV. Variants in the SP110 gene are associated with genetic susceptibility to tuberculosis in West Africa. *Proc Natl Acad Sci U S A.* 2006 Jul 5;103(27):10364-8. Epub 2006 Jun 27.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/16803959>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC1502463/>
- Watashi K, Hijikata M, Tagawa A, Doi T, Marusawa H, Shimotohno K. Modulation of retinoid signaling by a cytoplasmic viral protein via sequestration of Sp110b, a potent transcriptional corepressor of retinoic acid receptor, from the nucleus. *Mol Cell Biol.* 2003 Nov;23(21):7498-509.
Citation on PubMed: <https://www.ncbi.nlm.nih.gov/pubmed/14559998>
Free article on PubMed Central: <https://www.ncbi.nlm.nih.gov/pmc/articles/PMC207568/>

Reprinted from Genetics Home Reference:
<https://ghr.nlm.nih.gov/gene/SP110>

Reviewed: January 2009
Published: March 21, 2017

Lister Hill National Center for Biomedical Communications
U.S. National Library of Medicine
National Institutes of Health
Department of Health & Human Services